

T-odd Effects in Photon-Jet Production at the Tevatron

C. Pisano, D. Boer and P.J. Mulders

Vrije Universiteit Amsterdam, Department of Physics and Astronomy
NL-1081 HV Amsterdam, the Netherlands

The angular distribution in photon-jet production in $p\bar{p} \rightarrow \gamma \text{jet } X$ is studied within a generalized factorization scheme taking into account the transverse momentum of the partons in the initial hadrons. Within this scheme an anomalously large $\cos(2\phi)$ asymmetry observed in the Drell-Yan process could be attributed to the T-odd, spin and transverse momentum dependent parton distribution function $h_1^{\perp q}(x, \mathbf{p}_\perp^2)$. The same function is expected to produce a $\cos(2\phi)$ asymmetry in the photon-jet production cross section. This particular azimuthal asymmetry is estimated to be smaller than the Drell-Yan asymmetry but still of considerable size for Tevatron kinematics, offering a new possibility to study T-odd effects at the Tevatron.

In this contribution to DIS 2008 [1] we consider the process:

$$h_1(P_1) + h_2(P_2) \rightarrow \gamma(K_\gamma) + \text{jet}(K_j) + X, \quad (1)$$

where the four-momenta of the particles are given within brackets, and the photon-jet pair in the final state is almost back-to-back in the plane perpendicular to the direction of the incoming hadrons. To lowest order in pQCD the reaction is described in terms of the partonic two-to-two subprocesses $q(p_1) + \bar{q}(p_2) \rightarrow \gamma(K_\gamma) + g(K_j)$ and $q(p_1) + g(p_2) \rightarrow \gamma(K_\gamma) + q(K_j)$. Following reference [2], we will instead of collinear factorization consider a generalized factorization scheme taking into account partonic transverse momenta. We make a lightcone decomposition of the hadronic momenta in terms of two light-like Sudakov vectors n_+ and n_- , satisfying $n_+^2 = n_-^2 = 0$ and $n_+ \cdot n_- = 1$:

$$P_1^\mu = P_1^+ n_+^\mu + \frac{M_1^2}{2P_1^+} n_-^\mu, \quad \text{and} \quad P_2^\mu = \frac{M_2^2}{2P_2^-} n_+^\mu + P_2^- n_-^\mu. \quad (2)$$

In general n_+ and n_- will define the lightcone components of every vector a as $a^\pm \equiv a \cdot n_\mp$, while perpendicular vectors $a_\perp$ will always refer to the components of a orthogonal to both incoming hadronic momenta, P_1 and P_2 . Hence the partonic momenta (p_1, p_2) can be expressed in terms of the lightcone momentum fractions (x_1, x_2) and the intrinsic transverse momenta ($p_{1\perp}, p_{2\perp}$), as follows

$$p_1^\mu = x_1 P_1^+ n_+^\mu + \frac{m_1^2 + \mathbf{p}_{1\perp}^2}{2x_1 P_1^+} n_-^\mu + p_{1\perp}^\mu, \quad \text{and} \quad p_2^\mu = \frac{m_2^2 + \mathbf{p}_{2\perp}^2}{2x_2 P_2^-} n_+^\mu + x_2 P_2^- n_-^\mu + p_{2\perp}^\mu. \quad (3)$$

We denote with s the total energy squared in the hadronic center-of-mass (c.m.) frame, $s = (P_1 + P_2)^2 = E_{\text{c.m.}}^2$, and with η_i the pseudo-rapidities of the outgoing particles, *i.e.* $\eta_i = -\ln(\tan(\frac{1}{2}\theta_i))$, θ_i being the polar angles of the outgoing particles in the same frame. Finally, we introduce the partonic Mandelstam variables $\hat{s} = (p_1 + p_2)^2$, $\hat{t} = (p_1 - K_\gamma)^2$ and $\hat{u} = (p_1 - K_j)^2$, which satisfy the relations

$$-\frac{\hat{t}}{\hat{s}} \equiv y = \frac{1}{e^{\eta_\gamma - \eta_j} + 1}, \quad \text{and} \quad -\frac{\hat{u}}{\hat{s}} = 1 - y. \quad (4)$$




Figure 1: Cut diagrams for the subprocess $q\bar{q} \rightarrow \gamma g$.




Figure 2: Cut diagrams for the subprocess $qg \rightarrow \gamma q$.

Along the lines of [2, 3], we assume that at sufficiently high energies the hadronic cross section factorizes in a soft parton correlator for each observed hadron and a hard part:

$$d\sigma^{h_1 h_2 \rightarrow \gamma \text{jet} X} = \frac{1}{2s} \frac{d^3 K_\gamma}{(2\pi)^3 2E_\gamma} \frac{d^3 K_j}{(2\pi)^3 2E_j} \int dx_1 d^2 \mathbf{p}_{1\perp} dx_2 d^2 \mathbf{p}_{2\perp} (2\pi)^4 \delta^4(p_1 + p_2 - K_\gamma - K_j) \\ \times \sum_{a,b,c} \Phi_a(x_1, p_{1\perp}) \otimes \Phi_b(x_2, p_{2\perp}) \otimes |H_{ab \rightarrow \gamma c}(p_1, p_2, K_\gamma, K_j)|^2, \quad (5)$$

where the sum runs over all the incoming and outgoing partons taking part in the subprocesses $q\bar{q} \rightarrow \gamma g$ and $qg \rightarrow \gamma q$. The convolutions $\otimes$ indicate the appropriate traces over Dirac indices and $|H|^2$ is the hard partonic squared amplitude, obtained from the cut diagrams in Figs. 1 and 2. The parton correlators Φ_a describe the hadron $\rightarrow$ parton transitions and their parameterization in terms of transverse momentum dependent (TMD) distribution functions can be found, for instance, in [2].

We are interested in events in which the photon and the jet are approximately back-to-back in the transverse plane, therefore we define the vector $\mathbf{q}_\perp \equiv \mathbf{K}_{\gamma\perp} + \mathbf{K}_{j\perp}$ and require that $|\mathbf{q}_\perp| \ll |\mathbf{K}_{\gamma\perp}|, |\mathbf{K}_{j\perp}|$. Neglecting power-suppressed terms of the order $\mathcal{O}(1/(\mathbf{K}_{\gamma\perp}^4 s))$ and using the approximation $|\mathbf{K}_{\gamma\perp}| \approx |\mathbf{K}_{j\perp}|$, we obtain

$$\frac{d\sigma^{h_1 h_2 \rightarrow \gamma \text{jet} X}}{d\eta_\gamma d\eta_j d^2 \mathbf{K}_{\gamma\perp} d^2 \mathbf{q}_\perp} = \frac{1}{\pi^2} \frac{d\sigma^{h_1 h_2 \rightarrow \gamma \text{jet} X}}{d\eta_\gamma d\eta_j d\mathbf{K}_{\gamma\perp}^2 d\mathbf{q}_\perp^2} \left(1 + \mathcal{A}(y, x_1, x_2, \mathbf{q}_\perp^2) \cos 2(\phi_\perp - \phi_\gamma) \right), \quad (6)$$

with ϕ_γ and $\phi_\perp$ being the azimuthal angles, in the hadronic center-of-mass frame, of the outgoing photon and of the vector $\mathbf{q}_\perp$ respectively. The azimuthal asymmetry is given by

$$\mathcal{A}(y, x_1, x_2, \mathbf{q}_\perp^2) = \nu(x_1, x_2, \mathbf{q}_\perp^2) R(y, x_1, x_2, \mathbf{q}_\perp^2), \quad (7)$$

where ν contains the dependence on the time-reversal (T) odd distribution function $h_1^{\perp q}(x, \mathbf{p}_\perp^2)$ and is identical to the azimuthal asymmetry expression that appears in the Drell-Yan process [4], with the scale Q equal to $|\mathbf{K}_{\gamma\perp}|$. The function $h_1^{\perp q}(x, \mathbf{p}_\perp^2)$ is interpreted as the quark transverse spin distribution in an unpolarized hadron [5]. The role of gauge links in




Figure 3: The ratio R as a function of y , calculated according to (8) and (9) for different values of x_1, x_2 , $|\mathbf{K}_{\gamma\perp}|$ typical of the Tevatron experiments [6].

the calculation of ν is discussed in detail in [2]. The ratio R only depends on the T-even unpolarized distribution functions $f_1^{q,g}(x, \mathbf{p}_\perp^2)$, which integrated over $\mathbf{p}_\perp$ give the familiar lightcone momentum distributions $f_1^{q,g}(x)$. The explicit expressions of ν and R are given in [2].

The process $p\bar{p} \rightarrow \gamma \text{jet } X$ is currently being analyzed by the DØ Collaboration at the Tevatron collider. Data on the cross section, differential in η_γ, η_j and $\mathbf{K}_{j\perp}^2$, have been taken at $\sqrt{s} = 1.96$ TeV [6]. Such angular integrated measurements are only sensitive to the transverse momentum integrated parton distributions. A study of the angular dependent cross section in (6) will provide valuable information on the TMD distribution function $h_1^{\perp q}(x, \mathbf{p}_\perp^2)$, if the azimuthal asymmetry $\mathcal{A}$ turns out to be sufficiently sizeable in the available kinematic region. Model calculations [7, 8] applied to the $p\bar{p}$ Drell-Yan process have shown that the quantity ν in (7) is of the order of 30% or higher for $|\mathbf{q}_\perp|$ of a few GeV and Q values of $\mathcal{O}(1-10)$ GeV. Therefore, a study of the order of magnitude of $\mathcal{A}$ as a function of x_1, x_2 and $\mathbf{q}_\perp^2$ requires an estimate of the ratio R in (7). This will be obtained as follows.

First of all, the unknown TMD distribution functions appearing in the definition of R are evaluated assuming a factorization of their transverse momentum dependence, that is $f_1^{q,g}(x, \mathbf{p}_\perp^2) = f_1^{q,g}(x)\mathcal{T}(\mathbf{p}_\perp^2)$, with $f_1^{q,g}(x)$ being the usual unpolarized parton distributions and $\mathcal{T}(\mathbf{p}_\perp^2)$ being a generic function, taken to be the same for all partons and often chosen to be Gaussian. The $\mathbf{q}_\perp^2$ -dependence of R then drops out and we get

$$R = \frac{2N^2 y(1-y) \sum_q e_q^2 f_1^q(x_1) f_1^q(x_2)}{D(y, x_1, x_2)}, \quad (8)$$

where N is the number of colors and

$$D(y, x_1, x_2) = \sum_q e_q^2 \{ N(1-y)(1+y^2)f_1^q(x_1)f_1^q(x_2) + Ny(1+(1-y)^2)f_1^q(x_2)f_1^q(x_1) \\ + 2(N^2-1)(y^2+(1-y)^2)f_1^q(x_1)f_1^q(x_2) \}. \quad (9)$$

Here we have used that the antiquark contribution in the antiproton equals the quark contribution inside a proton. We consider only light quarks, *i.e.* the sum in (9) runs over $q = u, \bar{u}, d, \bar{d}, s, \bar{s}$, and we use the leading order GRV98 set [9] for the parton distributions, at the scale $\mu^2 = \mathbf{K}_{\gamma\perp}^2$.

Our results for R as a function of y are shown in Fig. 3 at some fixed values of the variables x_1, x_2 and $|\mathbf{K}_{\gamma\perp}|$, typical of the Tevatron experiments [6]. The values of x_1 and x_2 considered correspond to their average when both the photon and the jet are in the central rapidity region, where $\eta_j \approx \eta_\gamma \approx 0$ and $x_1 \approx x_2$. In this case $y \approx 0.5$, where R turns out to be largest. Evidently, R increases as x_1 and x_2 increase, due to the small contribution, in the denominator, of the gluon distributions $f_1^g(x)$ in the valence region.

Hence, we see that the asymmetry $\mathcal{A}$ is a product of a large Drell-Yan asymmetry term ν and a factor R that is estimated to be in the 10%-50% range for Tevatron kinematics. This leads us to conclude that an asymmetry $\mathcal{A}$ in the order of 5%-15% is possible in the central region. This could allow a study of the distribution function $h_1^{\perp q}$ in $p\bar{p} \rightarrow \gamma \text{jet } X$ at the Tevatron, offering a new possibility of measuring T-odd effects using a high energy collider.

Acknowledgments

We would like to thank C. Bomhof for useful comments and discussions. This research is part of the research program of the ‘‘Stichting voor Fundamenteel Onderzoek der Materie (FOM)’’, which is financially supported by the ‘‘Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO)’’.

References

- [1] Slides:
<http://indico.cern.ch/contributionDisplay.py?contribId=275&sessionId=22&confId=24657>
- [2] D. Boer, P. J. Mulders and C. Pisano, Phys. Lett. B **660**, 360 (2008).
- [3] A. Bacchetta, C. Bomhof, U. D’Alesio, P. J. Mulders and F. Murgia, Phys. Rev. Lett. **99** 212002 (2007).
- [4] D. Boer, Phys. Rev. D **60** 014012 (1999).
- [5] D. Boer and P. J. Mulders, Phys. Rev. D **57** 5780 (1998).
- [6] V. M. Abazov *et al.* [DØ Collaboration], arXiv:0804.1107 [hep-ex].
- [7] D. Boer, S. J. Brodsky and D. S. Hwang, Phys. Rev. D **67** 054003 (2003).
- [8] V. Barone, Z. Lu and B. Q. Ma, Eur. Phys. J. C **49** 967 (2007).
- [9] M. Glück, E. Reya and A. Vogt, Eur. Phys. J. C **5** 461 (1998).